

THE CONTRACT OF STATES OF THE STATES OF THE

AD-A192 815



OFFICE OF NAVAL RESEARCH

Contract N00014-83-K-0470-P00003

Task No. NR 359-718

TECHNICAL REPORT # 47

In Situ Infrared Spectroelectrochemistry

Ву

Stanley Pons C. Korzeniewski



Prepared for Publication in

Journal of Vacuum Science and Technology B

University of Utah
Department of Chemistry
Salt Lake City, Utah 84112

July 30, 1986

Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.

KITATAKA KITATA DALAK BANDA DALA

SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS HEFORE COMPLETING FORM
	2. GOVT ACCESSION NO.	3. HECIPIENT'S CATALOG NUMBER
47		
In Situ Infrared Spectroelectrocher	mistry	5. TYPE OF REPORT & PERIOD COVERED
•	·	Technical Report# 47
		6. PERFORMING ORG. REPORT NUMBER
7. ASTANTey Pons, C. Korzeniewski		B. CONTRACT OH GRANT NUMBER(s)
		N00014-83-K-0470_P0003
9. PERFORMING ORGANIZATION NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT HUMBERS
University of Utah Department of Chemistry	·	Task No. NR 359-718
Salt Lake City, UT 84112		
11. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE
Office of Naval Research	. A Co. d = . 470	July 30, 1986
Chemistry Program - Chemis Arlington, Virginia 22217	stry Lode 4/2	HUNDER OF FAUES
14. MONITORING AGENCY NAME & ADDRESS(II dillerent	from Controlling Office)	15. SECURITY CLASS. (of this report)
•	1	Unclassified
		ISA. DECLASSIFICATION/DOWNGRADING
		SCHEOULE
This document has been approved fo unlimited.	r public release	e and sale; its distribution
17. DISTRIBUTION STATEMENT (of the abstract entered in	Block 20, Il dillerent from	Report)
		•
	•	Į.
	•	
18. SUPPLEMENTARY NOTES		
		, · · · · · · · · · · · · · · · · · · ·
	•	·
	<u> </u>	
IR spectroelectrochemistry	identify by plock number)	ations
) 100	are. pur	
<b>\</b>		į
$\searrow$	•	į
The vibrational spectrum of molecular contractive vibrations of subtractively norm	dentily by block number)	an electrode surface can be
spectroscopy (SNIFTIRS) and electro		
, , , , , , , , , , , , , , , , , , , ,	•	
		Keynord

DD 10 1 1473 EDITION OF 1 HOV 65 IS ORSOLETE

Unclassified

AUTHOR PLEASE NOTE: All CorrectionS Must Be Marked On The Page Proof. Not On The Manuscript.

# XEROX COPY

MED PHYS, UVST, BAAS, HUMPORD, JOR PASTE 177 TO 100

010505JBS

Running Title: In situ infrared spectroelectrochemistry In situ infrared spectroelectrochemistry J. VAC. Sci. , TECH. B 3 (1985) 1421

Carol Korzeniewski and Stanley Ponsal Department of Chemistry. University of Utah, Salt Lake City, Utah 84112

(Received 15 February 1985; accepted 1 May 1985)

The vibrational spectrum of molecules at or near an electrode surface can be obtained using subtractively normalized interfacial Fourier transform infrared spectroscopy (SNIFTIRS) and electromodulated infrared spectroscopy (EMIRS). The large electric field existing at the electrode/solution interface is sufficient to induce dipole moments in highly polarizable molecules. The magnitude of the induced dipole moment is proportional to the electric field strength. As a result of this interaction some totally symmetric normal vibrations are made infrared active, and other bands appear in violation of the surface selection rule. This effect is known as the electrochemical Stark effect. Calculation of the expected absorption coefficients for the totally symmetric modes of adsorbed species gives values which are close to those observed experimentally. Experimental evidence of  $A_g$  mode activation of adsorbed pyrene is presented. The electric field dependence of the integrated absorption coefficient is demonstrated for acrylonitrile adsorbed on a gold electrode.

### I. INTRODUCTION

In recent years several techniques have been developed for obtaining in situ the infrared spectra of species at the electrode-solution interface. 1.2 The vibrational spectra is strongly dependent upon the molecular orientation with respect to the electrode surface. Electromagnetic radiation polarized perpendicular to the metal surface (s polarized) undergoes a phase shift close to 180° for all angles of incidence resulting in a standing wave which has little amplitude at the metal surface. The electric vectors of light polarized parallel to the plane of incidence (p polarized) add constructively upon reflection at glancing angles giving rise to a sizable component of electric vector normal to the surface. Therefore, only molecules which have a component of their dipole moment normal to the surface are able to interact with electromagnetic

Since intense electric fields exist near electrode surfaces, it is possible for the electric field to interact with highly polarizable molecules. This interaction may distort the diffuse electron cloud inducing a dipole in the direction of the electric field. If the polarization of the infrared radiation is in the same direction as the induced dipole, the integrated absorption coefficient for the transition is proportional to the square of the electric field strength. Therefore, the electric field at the electrode surface may allow infrared activity for vibrations normally forbidden by symmetry or the surface selection rule. This effect is known as the electrochemical Stark effect.<sup>6</sup> Perturbation of vibrational spectra by strong electric fields was first predicted by Condon<sup>3</sup> and has been demonstrated in several systems.

We report in this work a brief review and some new results that are related to the electrochemical Stark effect and vibronic activation of symmetric vibrational modes. The techniques used are modulated potential specular reflectance types that have recently been introduced to electrochemical science.


### II. EXPERIMENTAL

Experiments were performed in a three-electrode thinlayer cell (Fig. 1). The working electrode consisted of an 8-

10-mm-diam platinum or gold disk mounted on the end of a glass syringe barrel. The electrodes were polished to a mirror finish using alumina of decreasing sizes down to 0.05  $\mu$ m, then positioned in the cell so as to trap a thin layer of solution between the metal surface and the single crystal silicon infrared transmission window. The cell was then mounted in the path of a focused infrared light beam such that the angle of incidence from the normal was slightly less than 70 °C.

Spectra were obtained using both subtractively normalized Fourier transform infrared spectroscopy (SNIFTIRS) and electromodulated infrared spectroscopy (EMIRS)

In the standard SNIFTIRS technique, interferograms are collected and coadded at each of two electrode potentials. After transformation of the interferograms to the frequency domain spectra, the two results are ratioed. The final spectrum represents the difference in the infrared vibrational structure of the interfacial region at the two potentials. If transmission notation is chosen for the display of the spectrum, then the result is in units of  $1 + \Delta R / R$ , where  $\Delta R / R$ . the normalized reflectivity, is composed of the difference in reflectivity of the surface region at the two potentials, and R is the reflectance of the electrode at the potential corresponding to the state used in the denominator of the original



SCHEMATIC DIAGRAM OF ELECTROCHEMICAL INFRARED CELL

FIG. 1. Design of the thin layer spectroelectrochemical cell used for EMIRS and SNIFTIRS experiments.

0,0 05 103

Pg. No. Author: Title

ratio. At small values of  $\Delta R$  compared to R,  $\Delta R / R$  corresponds to the absorbance  $\Delta$ . + hrough put

In EMIRS, a high throughoun dispersive monochromator is used. The potential at the electrode is rapidly modulated between the potentials of interest, thereby modulating the relative amounts of absorbing species in the interfacial region. This results in a small modulation of the reflectivity which is easily extracted by synchronous demodulation techniques. Details of both experimental techniques have been presented earlier. 1.7

For cleaning cells, triply distilled water was used for steaming previously acid-cleaned cells. Aqueous solutions were prepared from recrystallized salts and triply distilled water. Acetonitrile (Burdick and Jackson, Muskegon MI) was used as received. The water content was nominally < 0.01%. Organic substrates were obtained from Aldrich and used as received. Electrochemistry was controlled by a potentiostat and waveform generator (JAS Instrument Systems, Inc.). Cells and electrodes were also obtained from JAS.

Typically, spectra were obtained at an electrode potential where the species of interest was either not adsorbed or weakly adsorbed; and at a potential where the species was present on the electrode surface or interfacial region to a large extent. The values of these potentials can be obtained usually by conventional double layer capacitance measurements which can give quantitative values of the surface coverage and information on the thickness of the electrical double layer near the interface.

### III. RESULTS AND DISCUSSION

### A. Calculation of B

In an experiment where the electric field  $(\vec{E}_x)$  is applied in a direction z, the induced dipole moment becomes


$$\mu_{\iota} = \alpha_{\iota \iota} \overline{E}_{\iota},$$

where  $\alpha_{x}$  is the polarizability tensor. The integrated absorption coefficient is given by

$$B = \frac{2\pi^2 vT}{\epsilon_0 hc} |\mu_{fi}|^2,$$

where  $\mu_A$  is the transition dipole matrix element. T is the number of absorbing molecules per unit area in the beam path, and all other constants have their usual meaning. In the presence of an external electric field the transition dipole matrix element can be expressed as the sum of the permanent

TABLE I.	Predicted values Son B. *						
	8/10 <sup>-1</sup> cm <sup>-1</sup> 54 Mode						
E V/m	C'H'	Napthalene Anthracene TCNQ			Pyrene		
10"		0.056	0.1	1.5	0.4		
10°	0.6	5.6	11	150	36		
2 < 10°	2 4	22	44	580	140		
5 × 10"	15	140	280	3 600	900		
8 × 10"	40	350	710	9 300	2 300		
10 "	62	560	1 100	14 500	3 600		



dipole moment ( $\mu_0$ ) and the induced dipole moment ( $\mu_z$ ) where

$$\mu_0 = \Sigma_q \langle r \rangle$$
.

Thus,

$$|\mu_A| = \langle \Psi_{\nu I} | \mu_0 + \mu_z | \Psi_0 \rangle_{-1}$$

If the expression for  $\mu_0$  and  $\mu_r$  are expanded in a Taylor series and the high order terms are neglected the integrated absorption coefficient becomes:

$$B = \frac{2\pi^2 vT}{\epsilon_0 hc} \left[ \langle \Psi_{i,j} | \mu_0^* | \Psi_{i,j} \rangle + E \langle \Psi_{i,j} | \alpha^* | \Psi_{i,j} \rangle \right]$$

where  $\mu_0'$  and  $\alpha'$  correspond to the change in the permanent dipole moment and polarizability with respect to a normal coordinate, respectively. We have calculated the absorption coefficient for the C=C symmetric stretch of molecules adsorbed flat at an electrode surface. For these molecules  $\mu_0$  approaches zero and we are left with evaluating the remaining matrix element. Since the matrix elements  $(\Psi, \alpha', \Psi, \gamma)$  are not tabulated for molecules larger than diatomic, we have estimated this change to be of the same order of magnitude as the polarizability normal to the molecular axis, providing an upper bound value to the matrix element. The polarizabilities necessary for the calculation were taken from Ref. 11. The value of the absorption coefficient is calculated as a function of electric field strength and shown in Table I.

### $A_{\rm r}$ mode activation

We have observed electric field activation of the totally symmetric  $A_e$  mode of pyrene at or near the surface of a platinum electrode. Figure 2 is the SNIFTIRS spectra of 0.5

Accession For

NTIS GRANT
DTIC TAB
Unannounced
Justification

By
Distribution/
Availability Codes

Avail and/or
Dist
Special

COPY NSPECTED

sa interes de Enorto de

010 525 712

TT CLE

MED PHYS. GVST. BAAS, AJ. LPCPD. JP PAGTERS IC 1300




FIG. 3. Raman spectrum of pyrene in the region 1485–1800 cm  $^{-1}$ . Ag mode assignments taken from Ref. 8.

mM pyrene in acetonitrile. The limits of the modulation potential were set between +0.5 and -1.2 V vs Ag/Ag\* where no electrochemical reaction was occurring. Therefore, the bands appearing in the difference spectra are due to the changing electric field rather than a chemical process. The band which occurs at  $\sim 1640$  cm $^{-1}$  corresponds to the totally symmetric  $(A_g)$  C-C stretching mode of pyrene. The Raman spectrum of pyrene was obtained in our labs and three bands were found in this frequency region (Fig.  $\bar{3}$ ). The





Fig. 4. EMIRS spectrum of 0.05 M acrylonitrile adsorbed at a gold electrode in 1 M H SO, as a function of potential vs SCE.






FIG. 5 Potential dependence of the intensity of the C=N stretch band from adsorbed acrylonitrile. The dashed curve is calculated from the squared electric field dependence.

two bands a (1628 and 1644 cm<sup>-1</sup> are assigned to the A-C-C stretching mode of pyrene. The positions of the activated band is very dependent on the potential, and is the subject of a forthcoming report. 10

### C. Electric field dependence of B

An example of the field dependence of band intensity of an adsorbed oscillator is demonstrated in the acrylonitrile system. Differential capacity measurements show that acrylonitrile is adsorbed on gold over a wide range of potentials (  $\pm$  0.5 $\pm$  1.0 V vs SCE). Spectroscopic evidence indicates that the molecule is adsorbed flat and interacts with the surface via the conjugated  $\pi$  system.

Figure 4 is the EMIRS spectra of acrylonitrile adsorbed on gold. The band at about 2130 cm<sup>-1</sup> corresponds to the C=N stretching mode of adsorbed acrylonitrile. The intensity of the band is about an order of magnitude weaker than expected for perpendicular adsorption, and is strongly potential dependent.7 As the potential is made more positive the band intensity increases. Figure 5 shows the band intensity as a function of the amplitude of the potential modulation. If only desorption of acrylonitrile at higher potentials was occuring, the band intensity should increase over the entire potential range and a band of opposite sign corresponding to solution free acrylonitrile should appear. However, as the modulation amplitude increases the band intensity increases, reaches a maximum, and begins to decrease. The observed spectra can be explained in terms of an electric field effect.

Since the potential dependence of the electric field at the electrode surface is unknown, it is difficult to test the exact field dependence of the intensity. However, there should be a maximum in band intensity when one of the modulation limits is at the potential point of zero charge at the interface. The dashed curve sketched in the figure is the theoretical curve predicted by the squared electric field dependence of intensity. The maximum in the curve appears at about -0.7 V/s SCE which is not an unreasonable position for the potential of zero field.

Also, the position of the band is not a linear function of

PROGRAEAU NG

010 555 513

TRHYS. CIIST. BAAS, AJ. JPCRD. JCP PASTER P. 104 700

Author: Title Pg. No.

potential. In systems were perpendicular adsorption is certain, such as CO adsorbed on platinum and CN adsorbed on silver, band position is a linear function of potential.

Although the fit is only approximate it is sufficiently good to demonstrate the effect of strong electric fields on infrared spectra. Intense electric fields present near the electrode surface may interact with polarizable molecules and induce a dipole moment in the direction of the field which give rise to very weak absorptions. With refinements in the calculation it may be possible to use the electrochemical Stark effect to study the electric field intensity in the double layer region of the electrode.

\* To whom correspondence should be addressed. S. Pons, J. Electroanal. Chem. 150, 495 (1983). -- INDENT

A. Bewick, K. Kunimatsu, B. S. Pons, and J. W. Russell, J. Electron-nal. Chem. 160, 47 (1984).

E. U. Condon, Phys. Rev. 41, 759 (1932).

M. F. Crawford and R. E. MacDonald, Can J. Phys. 36, 1022 1958. E. D. Palik, R. T. Holm, A. Stella, and H. L. Hughes, J. Applied Phys. 53, £454 (1987)

C. Korzeniewski, R. B. Shirts, and S. Pons, I. Phys. Chem. un pressi. A. Bewick and S. Pons, in Advances in Infrared and Raman Spectroscopy edited by R. Hester and R. Clark (Hayden, London, 1985)

A. Bree, R. A. Kydd, T. N. Missa, V. V. B. Vilkos, Spectrochim. Acta A. R7, 2315 (1971).

A. Bewick, C. Gibilaro, and S. Pons, Langmuir submitted C. Korzeniewski and S. Pons iin preparationi

"C.H., Naphlene, Anthracene: A. Chablo and A. Hinchliffe, Chem. Phys. Lett. 72, 149 (1980).

TCNQ": V. E. Klimenko, Ukr. Fiz. Zh. (Russ. Ed 123, 1301-1978 . CA) 89: 172337u.

Pyrene: J. Waite, M. G. Papaduopoulos, and C. A. Nicolaides, J. Chem. Phys. 77, 2536 (1982).

- : IDENT

09, 2297 (1986).

197 07 7717

# TECHNICAL REPORT DISTRIBUTION LIST, GEN

	No. Copies		No. Copies
Office of Naval Research Attn: Code 413 800 N. Quincy Street Arlington, Virginia 22217	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 5042 Crane, Indiana 47522	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Naval Civil Engineering.Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 2770	) 1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	r 12	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 1911	1
DTNSRDC Attn: Dr. G. Bosmajian Applied Chemistry Division Annapolis, Maryland 21401	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
Dr. William Toll <b>es</b> Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1		

Dr. Paul Delahay Department of Chemistry New York University New York, New York 10003

Dr. P. J. Hendra
Department of Chemistry
University of Southampton
Southampton S09 5NH
United Kingdom

Dr. J. Driscoll Lockheed Palo Alto Research Laboratory 3251 Hanover Street Palo Alto, California 94304

Dr. D. N. Bennion
Department of Chemical Engineering
Brigham Young University
Provo, Utah 84602

Dr. R. A. Marcus Department of Chemistry California Institute of Technology Pasadena, California 91125

Dr. J. J. Auborn
Bell Laboratories
Murray Hill, New Jersey 07974

Dr. Joseph Singer, Code 302-1 NASA-Lewis 21000 Brookpark Road Cleveland, Ohio 44135

Or. P. P. Schwift
Department of Chemistry
Oakland University
Rochester, Michigan 48063

Or. Manfred Breiter
Institut fur Technische Elektrochemie
Technischen Universitat Wien
9 Getreidemarkt, 1160 Wien
AUSTRIA

Dr. E. Yeager Department of Chemistry Case Western Reserve University Cleveland. Ohio 44106

Dr. C. E. Mueller
The Electrochemistry Branch
Naval Surface Weapons Center
White Oak Laboratory
Silver Spring, Maryland 20910

Dr. Sam Perone
Chemistry & Materials
Science Department
Lawrence Livermore National Laboratory
Livermore, California 94550

Dr. Royce W. Murray Department of Chemistry University of North Carolina Chapel Hill, North Carolina 27514

Dr. B. Brummer EIC Incorporated 111 Downey Street Norwood, Massachusetts 02062

Dr. Adam Heller Bell Laboratories Murray Hill, New Jersey 07974

Dr. A. B. Ellis Chemistry Department University of Wisconsin Madison, Wisconsin 53706

Library
Duracell, Inc.
Burlington, Massachusetts 01803

Electrochimica Corporation 20 Kelly Court Menlo Park, California 94025-1418

Dr. M. Wrighton
Chemistry Department
Massachusetts Institute
of Technology
Cambridge, Massachusetts 02139

Or. B. Stanley Pons
Department of Chemistry
University of Utah
Salt Lake City, Utah 84112

Donald E. Mains
Naval Weapons Support Center
Electrochemical Power Sources Division
Crane, Indiana 47522

S. Ruby
DOE (STOR)
Room 5E036 Forrestal Bldg., CE-14
Washington, D.C. 20595

Dr. A. J. Bard
Department of Chemistry
University of Texas
Austin, Texas 78712

Or. Janet Osteryoung
Department of Chemistry
State University of New York
Buffalo, New York 14214

Dr. Donald W. Ernst Naval Surface Weapons Center Code R-33 White Oak Laboratory Silver Spring, Maryland 20910

Mr. James R. Moden.
Naval Underwater: Systems Center
Code 3632
Newport, Rhode Island 02840

Dr. Bernard Spielvogel
U.S. Army Research Office
P.O. Box 12211
Research Triangle Park, NC 27709

Dr. Aaron Fletcher Naval Weapons Center Code 3852 China Lake, California 93555 Dr. M. M. Nicholson Electronics Research Center Rockwell International 3370 Miraloma Avenue Anaheim, California

Dr. Michael J. Weaver Department of Chemistry Purdue University West Lafayette, Indiana 47907

Dr. R. David Rauh EIC Laboratories, Inc. 111 Downey Street Norwood, Massachusetts 02062

Dr. Aaron Wold Department of Chemistry Brown University Providence, Rhode Island 02192

Dr. Martin Fleischmann Department of Chemistry University of Southampton Southampton SO9 5NH ENGLAND

Dr. R. A. Osteryoung
Department of Chemistry
State University of New York
Buffalo, New York 14214

Dr. John Wilkes
Air Force Office of Scientific
Research
Bolling AFB
Washington, D.C. 20332

Dr. R. Nowak Naval Research Laboratory Code 6171 Washington, D.C. 20375

Dr. D. F. Shriver
Department of Chemistry
Northwestern University
Evanston, Illinois 60201

Dr. Hector D. Abruna
Department of Chemistry
Cornell University
Ithaca, New York 14853

Dr. A. B. P. Lever Chemistry Department York University Downsview, Ontario M3J1P3

Dr. Stanislaw Szpak Naval Ocean Systems Center Code 633, Bayside San Diego, California 95152

Dr. Gregory Farrington
Department of Materials Science
and Engineering
University of Pennsylvania
Philadelphia, Pennsylvania 19104

M. L. Robertson
Manager, Electrochemical
and Power Sources Division
Naval Weapons Support Center
Crane, Indiana 47522

Dr. T. Marks Department of Chemistry Northwestern University Evanston, Illinois 60201

Dr. Micha Tomkiewicz Department of Physics Brooklyn College Brooklyn, New York 11210

Dr. Lesser Blum
Department of Physics
University of Puerto Rico
Rio Piedras. Puerto Rico 00931

Dr. Joseph Gordon, II IBM Corporation 5600 Cottle Road San Jose, California 95193

Dr. Nathan Lewis Department of Chemistry Stanford University Stanford, California 94305 Dr. D. H. Whitmore
Department of Materials Science
Northwestern University
Evanston, Illinois 60201

Dr. Alan Bewick Department of Chemistry The University of Southampton Southampton, SO9 5NH ENGLAND

Dr. E. Anderson NAVSEA-56Z33 NC #4 2541 Jefferson Davis Highway Arlington, Virginia 20362

Dr. Bruce Dunn
Department of Engineering &
Applied Science
University of California
Los Angeles, California 90024

Dr. Elton Cairns
Energy & Environment Division
Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

Dr. Richard Pollard
Department of Chemical Engineering
University of Houston
Houston, Texas 77004

Dr. M. Philpott IBM Corporation 5600 Cottle Road San Jose, California 95193

Dr. Donald Sandstrom Boeing Aerospace Co. P.O. Box 3999 Seattle, Washington 98124

Dr. Carl Kannewurf
Department of Electrical Engineering
and Computer Science
Northwestern University
Evanston, Illinois 60201

Dr. Joel Harris Department of Chemistry University of Utah Salt Lake City, Utah 84112

Dr. Robert Somoano Jet Propulsion Laboratory California Institute of Technology Pasadena. California 91103

Dr. Johann A. Joebstl
USA Mobility Equipment R&D Command
DRDME-EC
Fort Belvoir, Virginia 22060

Dr. Judith H. Ambrus NASA Headquarters M.S. RTS-6 Washington, D.C. 20546

Dr. Albert R. Landgrebe U.S. Department of Energy M.S. 68025 Forrestal Building Washington, D.C. 20595

Dr. J. J. Brophy
Department of Physics
University of Utah
Salt Lake City, Utah 84112

Dr. Charles Martin
Department of Chemistry
Texas A&M University
College Station, Texas 77843

Dr. H. Tachikawa Department of Chemistry Jackson State University Jackson, Mississippi 39217

Dr. Theodore Back Electrochemical Technology Corp. 3935 Leary Way N.W. Seattle, Washington 98107

Dr. Farrell Lytle
Boeing Engineering and
Construction Engineers
P.O. Box 3707
Seattle, Washington 98124

Dr. Robert Gotscholl U.S. Department of Energy MS G-226 Washington, D.C. 20545 Dr. Edward Fletcher Department of Mechanical Engineering University of Minnesota Minneapolis, Minnesota 55455

Dr. John Fontanella Department of Physics U.S. Naval Academy Annapolis, Maryland 21402

Dr. Martha Greenblatt
Department of Chemistry
Rutgers University
New Brunswick, New Jersey 08903

Dr. John Wasson Syntheco, Inc. Rte 6 - Industrial Pike Road Gastonia, North Carolina 28052

Or. Walter Roth
Department of Physics
State University of New York
Albany, New York 12222

Dr. Anthony Sammells Eltron Research Inc. 4260 Westbrook Drive, Suite 111 Aurora, Illinois 60505

Dr. C. A. Angell Department of Chemistry Purdue University West Lafayette, Indiana 47907

Dr. Thomas Davis
Polymer Science and Standards
Division
National Bureau of Standards
Washington, D.C. 20234

Ms. Wendy Parkhurst
Naval Surface Weapons Center R-33
R-33
Silver Spring, Maryland 20910

Dr. John Owen
Department of Chemistry and
Applied Chemistry
University of Salford
Salford M5 4WT ENGLAND

Dr. Boone Owens
Department of Chemical Engineering
and Materials Science
University of Minnesota
Minneapolis, Minnesota 55455

Dr. J. O. Thomas University of Uppsala Institute of Chemistry Box 531 S-751 21 Uppsala, Sweden Dr. O. Stafsudd
Department of Electrical Engineering
University of California
Los Angeles, California 90024

Dr. S. G. Greenbaum Department of Physics Hunter College of CUNY New York, New York 10021

Or. Menahem Anderman W.R. Grace & Co. Columbia, Maryland 20144

IE DA